

Cool Orchard Temperatures or Growing Trees in Containers Can Inhibit Leaf Gas Exchange of Avocado and Mango

Anthony W. Whiley

Maroochy Research Station, Centre for Subtropical Fruit, Queensland Horticulture Institute, Department of Primary Industries, P.O. Box 5083 S.C.M.C., Nambour 4560, Australia

Christopher Searle

Bundaberg Research Station, Queensland Horticulture Institute, Department of Primary Industries, Ashfield Road, Kalkie 4670, Australia

Bruce Schaffer¹

Tropical Research and Education Center, Institute of Food and Agricultural Sciences, University of Florida, 18905 S.W. 280 Street, Homestead, FL 33031

B. Nigel Wolstenholme

Department of Horticultural Science, University of Natal, Private Bag X01 Scottsville, Pietermaritzburg 3209, South Africa

ADDITIONAL INDEX WORDS. *Persea americana*, *Mangifera indica*, chlorophyll fluorescence, photoinhibition, photosynthesis, photosynthetic photon flux

ABSTRACT. Leaf gas exchange of avocado (*Persea americana* Mill.) and mango (*Mangifera indica* L.) trees in containers and in an orchard (field-grown trees) was measured over a range of photosynthetic photon fluxes (PPF) and ambient CO₂ concentrations (C_a). Net CO₂ assimilation (A) and intercellular partial pressure of CO₂ (C_i) were determined for all trees in early autumn (noncold-stressed leaves) when minimum daily temperatures were ≥14 °C, and for field-grown trees in winter (cold-stressed leaves) when minimum daily temperatures were ≤10 °C. Cold-stressed trees of both species had lower maximum CO₂ assimilation rates (A_{max}), light saturation points (Q_A), CO₂ saturation points (C_{aSAT}) and quantum yields than leaves of noncold-stressed, field-grown trees. The ratio of variable to maximum fluorescence (F_v/F_m) was ≈50% lower for leaves of cold-stressed, field-grown trees than for leaves of nonstressed, field-grown trees, indicating chill-induced photoinhibition of leaves had occurred in winter. The data indicate that chill-induced photoinhibition of A and/or sink limitations caused by root restriction in container-grown trees can limit carbon assimilation in avocado and mango trees.

Root restriction from growing plants in containers can limit net CO₂ assimilation rates (A) via feedback inhibition (Arp, 1991; Schaffer et al., 1996; Thomas and Strain, 1991). For avocado trees, maximum A (A_{max}) ranged from ≈7 to 23 μmol·m⁻²·s⁻¹ with light saturation for A (Q_A) at photosynthetic photon fluxes (PPF) between 400 to 1100 μmol·m⁻²·s⁻¹ (Bower et al., 1978; Scholefield et al., 1980; Schaffer et al., 1987; Whiley, 1994; Whiley and Schaffer, 1994). Mango A_{max} was ≈6 to 18 μmol·m⁻²·s⁻¹ with Q_A at PPF between 300 to 1200 μmol·m⁻²·s⁻¹ (Chacko et al., 1995; Pongsomboon et al., 1992; Schaffer and Gaye, 1989; Schaffer et al., 1994; Searle et al., 1995). The range of variation in these data was possibly due to measurement or growth conditions, particularly since the lower values were from trees grown in containers, whereas higher values were from field-grown trees.

Received for publication 25 Mar. 1998. Accepted for publication 8 Sept. 1998. Financial support for this study was provided, in part, by the Rural Industries Research and Development Corporation (Australia) under the Climate Change Program. We thank V. Doogan of Biometry, Queensland Department of Primary Industries for assistance with curve fitting and T.S. Rasmussen of Agricultural Chemistry, Queensland Department of Natural Resources for carrying out the starch analyses. B. Schaffer thanks the Rural Industries Research and Development Corporation and the Queensland Department of Primary Industries for financial support while on study-leave at Maroochy Research Station, Centre for Subtropical Fruit, Queensland Horticulture Institute, Nambour, Australia. This article is Florida Agricultural Experiment Station journal series no. R-06241. The cost of publishing this paper was defrayed in part by the payment of page charges. Under postal regulations, this paper therefore must be hereby marked *advertisement* solely to indicate this fact.

¹To whom reprint requests should be addressed; e-mail: Bas@gvn.ifas.ufl.edu.

Short-term exposure to temperatures below 10 °C can cause reversible chilling injury resulting in inhibition of A, especially in subtropical and tropical species (Taylor and Rowley, 1971). At high incident PPF, chilling can result in photoinhibitory damage to photosystem 2 (PS II) (Powles, 1984; Smillie et al., 1988) that can be quantified by measuring a decrease in the ratio of variable to maximum chlorophyll fluorescence (F_v/F_m) (Björkman, 1987; Demmig and Björkman, 1987).

Avocado (*Persea americana* Mill.), a polyaxial, terminally flowering, oil-accumulating species (Whiley and Schaffer, 1994) and mango (*Mangifera indica* L.), a polyaxial, terminally flowering, sugar-accumulating species (Schaffer et al., 1994), are gaining commercial importance worldwide and production is expanding into new environments. Whiley (1994) has classified avocado as a “wintergreen” tree, which more closely resembles the physiology of woody deciduous species. There is comparatively little carbon investment into avocado leaves, which grow rapidly but have a longevity of <10 months. In contrast, mango leaves are relatively sclerophyllous with a high investment of carbon during growth. Furthermore, they have drought-resistant mechanisms and are sustained on trees for up to 5 years (Whiley and Schaffer, 1994). Quantifying leaf gas exchange characteristics of these similar evergreen, broad-leaved species may enable the development of models of plant responses to growth conditions (e.g., variation in temperatures and elevated ambient CO₂ concentrations), thereby minimizing deleterious effects from suboptimal environments (Schaffer and Andersen, 1994).

The purpose of this study was to compare photoinhibitory and sink-limitation (root restriction) responses between wintergreen (avocado) and sclerophyllous (mango) leaves of two broad-leaved evergreen trees. To assess potential photoinhibitory damage to leaves, A of field-grown trees was determined when minimum daily (overnight) temperatures were below 10 °C. Root restriction responses were assessed by comparing leaf gas exchange responses of field-grown and container-grown trees.

Materials and Methods

PLANT MATERIAL. Four- to six-year-old 'Hass' avocado and 'Kensington' (syn. 'Kensington Pride') mango trees were used in this study. Experiments were conducted with field- and container-grown (55-L black polythene containers) trees at Maroochy Research Station, Centre for Subtropical Fruits, Nambour, Queensland, Australia (lat. = 27° S, elevation = 30 m above sea level). The orchard soil was a well-drained clay loam (≈60% clay fraction) 10 to 16 m deep and showed no obvious physical limitation to root growth (Whiley, 1994). Trees in containers were grown outdoors under ambient conditions in a potting media of course river sand and peat (1:1). For each species, five single-plant replications were used for leaf gas exchange and leaf starch determinations in studies with both field- and container-grown trees. Plants were irrigated and fertilized according to standard commercial practices (Whiley, 1984; Whiley et al., 1988).

LEAF GAS EXCHANGE. Net CO₂ assimilation of leaves and intercellular partial pressure of CO₂ (C_i) were determined for the second or third most recently fully expanded leaf of field- and container-grown plants under a range of PPF and ambient CO₂ concentrations (C_a). Before measurements, plants were grown under full sunlight (maximum PPF of ≈2000 μmol·m⁻²·s⁻¹) and ambient atmospheric CO₂ concentrations (≈350 μmol·mol⁻¹). Determinations of A and C_i were made with a CIRAS-1 leaf gas exchange system (PP Systems, Hitchin, Herts., U.K.). Leaf gas exchange was measured when shoot growth on trees was quiescent during early autumn 1994 and minimum daily temperatures were ≥14 °C (noncold-stressed leaves = NCS), and again on field-grown trees in winter 1994 when minimum daily (overnight) temperatures were ≤10 °C (cold-stressed leaves = CS). Cold-stressed leaves were exposed to 24 nights of temperatures of 8 to 10 °C. Measurements in autumn were made between 0800 to 1030 HR when air temperatures were 25 to 28 °C, whereas winter measurements were between 0900 to 1130 HR when air temperatures were 21 to 24 °C. At each time of the year, vapor pressure deficits were 1.0 to 1.2 kPa when measurements were taken.

To investigate leaf gas exchange responses to incident light flux, PPF was varied nonconsecutively by placing 1 m² frames covered with polyethylene cloths of different mesh densities, and in different combinations, between the trees and the sun (on a cloudless day) to obtain a range of PPF. The entire canopy was shaded. Frames were placed over the trees for at least 30 min before A determinations to allow sufficient time for leaves to equilibrate to the new light environment. Determinations of A at zero PPF were obtained by wrapping the leaf cuvette in a black cloth and waiting until CO₂ evolution from the leaf stabilized (8 to 12 min).

Leaf gas exchange responses to C_a were made at saturating PPF (previously determined as >400 μmol·m⁻²·s⁻¹). A range of C_a concentrations in the leaf cuvette was achieved using the CIRAS-1 gas exchange system, which allows variable CO₂ concentrations to be delivered to the cuvette from a supplemented CO₂ source. For below ambient concentrations, the air stream was passed through a series of CO₂ scrubbers (cylinders containing soda lime) before

entering the cuvette. Determinations of A and C_i were made when CO₂ flux in the cuvette had equilibrated (≈3 to 5 min after placing the leaf in the cuvette after C_a was changed).

CHLOROPHYLL FLUORESCENCE. To quantify chilling injury to PS II, chlorophyll fluorescence of leaves on field-grown trees was measured on four to five leaves per tree at the same time as leaf gas exchange determinations were made using a BioMonitor Stress Meter (BioMonitor SCI, Umeå, Sweden). Cuvettes were attached to each side of the leaf midrib for 30 min between 0900 and 1000 HR. Chlorophyll fluorescence was then determined on the adaxial leaf surface after two seconds of irradiation with 600 μmol·m⁻²·s⁻¹ of blue light (320 to 550 nm). Variable fluorescence (F_v) was calculated as $F_v = F_m - F_o$, where F_o is the initial constant yield fluorescence and F_m is the maximum fluorescence recorded (Öquist and Wass, 1988). The ratio of F_v/F_m was calculated as an indication of photoinhibitory damage to PS II (Björkman, 1987; Demmig and Björkman, 1987).

LEAF STARCH. Starch concentrations in avocado and mango trees were determined on the second or third, fully expanded leaf from the most recently matured shoots on each tree. Fifteen leaves were collected from each of five trees from the different treatments between 0800 and 1030 HR and subsequently placed in a convection oven at 60 °C and dried to constant mass. Dried samples were ground at 100 mesh in a cyclone grinder (UDY Corporation, Fort Collins, Colo.) and stored in air-tight containers. Starch was determined by a two-stage enzymatic hydrolysis of starch to glucose and the concentration measured colorimetrically with a coupled glucose oxidase/oxidase/chromogen system as described by Rasmussen and Henry (1990).

DATA ANALYSES. Data were analyzed by regression analysis (Table Curve, SPSS, Inc., Chicago). Light and CO₂ saturation points for A were calculated as 90% of A_{max} (Osman and Milthorpe, 1971). Quantum yield (Φ) was calculated as the slope of the linear portion of the regression line of A vs. incident PPF (Syvertsen, 1984). Significant differences in Φ between field- and container-grown plants of each species were determined by testing for homogeneity of slopes (SAS Institute, Cary, N.C.).

Results

LEAF GAS EXCHANGE RESPONSES TO INCIDENT PPF. Net CO₂ assimilation increased and C_i decreased asymptotically in response to increasing PPF for both species (Figs. 1 and 2). For both species, A_{max} was greater for field- than for container-grown trees and greater for NCS than for CS leaves (Table 1). For avocado, the light saturation point for A (Q_A) was ≈1270 μmol·m⁻²·s⁻¹ for NCS, field-grown trees, compared to ≈1040 μmol·m⁻²·s⁻¹ for CS, field-grown trees and ≈586 μmol·m⁻²·s⁻¹ for container-grown trees (Table 1). The light compensation point (Q_o) for field-grown avocado was lower for NCS (30 μmol·m⁻²·s⁻¹) than for CS (50 μmol·m⁻²·s⁻¹) trees. For container-grown trees, Q_o was 38 μmol·m⁻²·s⁻¹. Cold stress reduced Φ of field-grown trees from 0.055 μmol CO₂/mol quanta in fall to 0.034 μmol CO₂/mol quanta following exposure to overnight minimums of ≤10.0 °C, whereas the Φ for container-grown trees was further reduced to 0.021 μmol CO₂/mol quanta. Quantum yield was higher in NCS than in CS leaves ($P \leq 0.01$) and in field- compared with container-grown avocado trees ($P \leq 0.05$).

For container- and field-grown avocado trees, C_i initially declined rapidly and then leveled off at PPF above 500 μmol·m⁻²·s⁻¹ (Fig. 1b). To quantify the effects of PPF on C_i , the $Q_{1/2}$, representing the PPF at which C_i is reduced by 50%, was calculated from $Q_{1/2} = \ln(0.5)/-k$, where k is the third constant in the standard decay curve fitted to the data (adapted from Jones, 1992).




Fig. 1. Net CO₂ assimilation (*A*) and intercellular partial pressure of CO₂ (*C_i*) responses of field- and container-grown avocado trees ('Hass') to varying photosynthetic photon fluxes (PPF). The regression line for *A* (a) of noncold-stressed (NCS), field-grown trees is represented by $y = 22.08(-30.67 + x)/(427.43 + x)$, $r^2 = 0.94$; cold stressed (CS), field-grown trees is represented by $y = 14.17(-46.72 + x)/(250.05 + x)$, $r^2 = 0.86$; and container-grown trees is represented by $y = 7.16(-38.366 + x)/(172.12 + x)$, $r^2 = 0.84$. The regression line for *C_i* (b) for NCS, field-grown trees is represented by $y = 244.65 + 90.2 \exp(-0.0034x)$, $r^2 = 0.94$; CS, field-grown trees by $y = 215.28 + 135.52 \exp(-0.0041x)$, $r^2 = 0.92$; and container-grown trees by $y = 203.70 + 171.39 \exp(-0.0056x)$, $r^2 = 0.87$.

The $Q_{1/2}$ for NCS, field-grown avocado trees was 204 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ compared to 169 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ for CS, field-grown trees and 123 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ for container-grown trees, indicating that leaves from CS, field-grown and container-grown trees had a higher light requirement than leaves of NCS, field-grown trees to reduce *C_i* by one-half.

For NCS, field-grown mango trees, Q_A was $\approx 1284 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$, compared to $\approx 1180 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ for CS, field-grown trees and $\approx 563 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ for container-grown trees (Table 1). The Q_o for field-grown mango was 29 and 66 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ for NCS and CS leaves, respectively. For container-grown trees, Q_o was 47 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$. Lower temperatures reduced Φ of field-grown trees from 0.042 $\mu\text{mol CO}_2/\mu\text{mol quanta}$ in fall to 0.025 $\mu\text{mol CO}_2/\mu\text{mol quanta}$ for CS leaves in winter, whereas the Φ for container-grown trees was 0.033 (Table 1). Quantum yield was significantly greater between NCS and CS leaves ($P \leq 0.01$) as well as between NCS, field- and container-grown mango trees ($P \leq 0.05$).

The response of mango *C_i* to PPF was similar to that of avocado trees: *C_i* decreased as PPF increased but leveled off as A_{max} was reached (Fig. 2b). The $Q_{1/2}$ for leaves from NCS, field-grown mango trees was 223 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ compared to 74 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ for CS, field- and 161 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ for container-grown trees.

LEAF GAS EXCHANGE RESPONSES TO AMBIENT PARTIAL PRESSURE OF CO₂. For container- and field-grown avocado and mango trees, *A* increased asymptotically as *C_a* increased, whereas there was a

linear relationship between *C_i* and *C_a* (Figs. 3 and 4). The ambient CO₂ concentration at which A_{max} occurred (C_{aSAT}) in avocado was highest for leaves from NCS, field-grown trees (1473 $\mu\text{mol}\cdot\text{mol}^{-1}$) compared with leaves from CS, field- (1357 $\mu\text{mol}\cdot\text{mol}^{-1}$) and container-grown (1261 $\mu\text{mol}\cdot\text{mol}^{-1}$) trees (Table 2). The saturated CO₂ assimilation rate (A_{maxCa}) was also lower for leaves from CS, field- ($A_{maxCa} = 34.9 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) and container-grown ($A_{maxCa} = 26.1 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) avocado trees than for NCS, field-grown avocado trees ($A_{maxCa} = 50.6 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) (Table 2).

For leaves of NCS, field-grown mango, C_{aSAT} was 1113 compared to 871 $\mu\text{mol}\cdot\text{mol}^{-1}$ for container-grown and 636 $\mu\text{mol}\cdot\text{mol}^{-1}$ for CS, field-grown trees (Table 2). At *C_a* levels >600 $\mu\text{mol}\cdot\text{mol}^{-1}$, *A* tended to level off for field- and container-grown trees (Fig. 4a) while A_{maxCa} was higher for NCS, field- (30.3 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) than for container-grown (21.5 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) and CS, field-grown (8.3 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) mango trees (Table 2). For container- and field-grown avocado and mango trees, *C_i* increased linearly as *C_a* increased indicating that CO₂ concentration was not limiting *A* (Figs. 3b and 4b).

CHLOROPHYLL FLUORESCENCE. The F_v/F_m ratios for leaves of NCS, field-grown trees were 0.81 ± 0.02 and 0.80 ± 0.03 , for avocado and mango, respectively (Table 2). During winter, the F_v/F_m ratio declined to 0.41 ± 0.03 for avocado and 0.45 ± 0.03 for mango.

LEAF STARCH. Leaf starch concentrations (leaf dry weight basis) measured in NCS, field-grown avocados ($17.4 \pm 1.4 \text{ mg}\cdot\text{g}^{-1}$) were less than those for CS, field- ($23.4 \pm 4.0 \text{ mg}\cdot\text{g}^{-1}$) and container-




Fig. 2. Net CO₂ assimilation (*A*) and intercellular partial pressure of CO₂ (*C_i*) responses of field- and container-grown mango trees ('Kensington') to varying photosynthetic photon fluxes (PPF). The regression line for *A* (a) of noncold-stressed (NCS), field-grown trees is represented by $y = 20.94(-28.97 + x)/(448.27 + x)$, $r^2 = 0.94$; cold stressed (CS), field-grown trees by $y = 12.42(-66.26 + x)/(309.02 + x)$, $r^2 = 0.91$; and container-grown trees by $y = 11.40(-45.97 + x)/(152.39 + x)$, $r^2 = 0.91$. The regression line for *C_i* (b) for NCS, field-grown trees is represented by $y = 267.27 + 87.38 \exp(-0.0031x)$, $r^2 = 0.97$; for CS, field-grown trees by $y = 245.57 + 133.54 \exp(-0.0094x)$, $r^2 = 0.93$; and container-grown trees by $y = 255.0 + 90.46 \exp(-0.0043x)$, $r^2 = 0.95$.

Table 1. The effect of growing conditions on light saturation of CO₂ assimilation (Q_A), the compensation point for CO₂ assimilation (Q_o), the maximum rate of CO₂ assimilation (A_{max}) and the quantum efficiency (ϕ) of avocado and mango trees. Values were estimated from the regression models presented in Figs. 1 and 2, which were based on pooled data from one leaf of each of five trees.

Treatment	Q_A ($\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$)	Q_o ($\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$)	A_{max} ($\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$)	ϕ ($\mu\text{mol CO}_2/\mu\text{mol quanta}$)
Avocado				
NCS field-grown	1270	30	19.0	0.055a ^z
CS field-grown	1040	50	10.9	0.034b
Container-grown	586	38	5.2	0.021c
Mango				
NCS field-grown	1284	29	15.2	0.042a ^z
CS field-grown	1180	66	8.8	0.025b
Container-grown	563	47	8.1	0.033c

^zMeans followed by different letters for each species indicate significant difference ($P \leq 0.05$) according to a test for homogeneity of slopes.

grown trees ($28.0 \pm 2.6 \text{ mg}\cdot\text{g}^{-1}$) (Table 2). There was also a similar response for leaf starch concentrations in mango with levels in NCS, field-grown trees at $14.0 \pm 0.30 \text{ mg}\cdot\text{g}^{-1}$ dry weight compared with $25.4 \pm 1.4 \text{ mg}\cdot\text{g}^{-1}$ for leaves of CS, field-grown and $32.2 \pm 1.8 \text{ mg}\cdot\text{g}^{-1}$ for container-grown trees.

Discussion

The lower A of container-grown than of field-grown plants may have been due to a carbon sink limitation as a result of root restriction in containers. Although the root mass was not determined for container- and field-grown plants in this study, previous experiments have indicated that similar-aged avocado and mango plants maintained in similar-size containers for the time period of this study become pot bound (A.W. Whiley and B. Schaffer, unpublished data). Cold stress (exposing trees to 24 d of 8 to 10 °C temperatures) also limited A of avocado and mango in this study. Below saturating PPF, C_i decreased linearly as PPF increased, indicating unrestricted CO₂ diffusion through stomata. Thus, the reductions in A were apparently due to reduced carboxylation by the plant rather than a reduction in CO₂ diffusion through the mesophyll tissues. However, at Q_A , C_i remained relatively constant indicating a point of equilibrium between PPF and the diffusion of CO₂ to fixation sites. Under noncold-stressed conditions, ϕ for avocado and mango trees approximated the normal range of ϕ for C₃ species (Ehleringer and Björkman, 1977). The lower ϕ for CS, field- and container-grown trees than for NCS trees indicated a reduced quantum-use efficiency at low incident PPF as a result of both source and sink limitations to A .

The Q_A observed in this study for NCS, field-grown avocado ($Q_A \approx 1270 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) and mango ($Q_A \approx 1284 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) were considerably higher than previously reported rates for these crops (Bower et al., 1978; Scholefield et al., 1980; Schaffer and Gaye, 1989). The higher $Q_{1/2}$ for NCS, field-grown avocado trees than for the CS, field-grown trees and container-grown trees, indicates that leaves of CS, field-grown and container-grown trees had a higher light requirement than leaves of NCS, field-grown trees to reduce C_i by one-half. Thus, the Q_A of NCS leaves would be higher than that of leaves from CS, field- and container-grown trees (assuming stomatal conductance is nonlimiting under well-watered conditions), thereby supporting the data presented in Fig. 1. The A_{max} that we observed for NCS, field-grown trees of each species was much greater than data previously published. For field-grown avocado in Florida, A_{max} was reported to be 7 to 10 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ (Schaffer et al., 1987, 1991) compared to 19.0 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ observed for field-grown trees in this study. Similarly, A_{max} for field-grown mango in Florida was 6.2 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ (Schaffer and Gaye, 1989) compared

to 15.2 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ determined in our study for nonstressed field-grown trees in Australia. In contrast, the A_{max} and Q_A for container-grown avocado (5.2 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$; 586 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) and mango (8.1 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$; 563 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) were similar to those previously reported for trees grown in containers (Bower et al., 1978; Scholefield et al., 1980; Schaffer et al., 1987; Schaffer and Gaye, 1989; Larson et al., 1992; Pongsomboon et al., 1992). For single leaves of container-grown avocado, Bower et al. (1978) and Scholefield et al. (1980) have reported Q_A to be between 400 to 660 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ with an A_{max} of $\approx 7.0 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$, while for mango a Q_A of 350 to 400 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ with an A_{max} of 6 to 7 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ has




Fig. 3. Net CO₂ assimilation (A) and intercellular partial pressure of CO₂ (C_i) responses of field- and container-grown avocado ('Hass') trees to varying ambient CO₂ partial pressures (C_a). The regression line for A of noncold-stressed (NCS), field-grown trees (a) is represented by $y = 85.49((-96.72 + x)/(852.75 + x))$, $r^2 = 0.96$; cold-stressed (CS), field-grown trees by $y = 58.54((-110.10 + x)/(805.77 + x))$, $r^2 = 0.96$; and container-grown trees by $y = 42.73((-78.62 + x)/(801.42 + x))$, $r^2 = 0.92$. The regression line for C_i (b) of NCS, field-grown trees is represented by $y = -0.851x - 47.590$, $r^2 = 0.99$; for CS, field-grown trees by $y = 0.812x - 60.877$, $r^2 = 0.98$; and container-grown trees by $y = 0.723x - 27.95$, $r^2 = 0.95$.




Fig. 4. Net CO₂ assimilation (*A*) and intercellular partial pressure of CO₂ (*C_i*) responses of field- and container-grown mango ('Kensington') trees to varying ambient CO₂ partial pressures (*C_a*). The regression line for *A* (a) for noncold-stressed (NCS), field-grown trees is represented by $y = 47.38((-105.0 + x)/(462.4 + x))$, $r^2 = 0.95$; cold stressed (CS), field-grown trees by $y = 10.24 - 1247.52/x$, $r^2 = 0.73$; and container-grown trees by $y = 33.61 - 357.84/x^{0.5}$, $r^2 = 0.91$. The regression line for *C_i* (b) for NCS, field-grown trees is represented by $y = 9.026 + 0.753x$, $r^2 = 0.99$; CS, field-grown trees by $y = 40.67 + 0.568x$, $r^2 = 0.92$; and container-grown trees by $y = -10.857 + 0.872x$, $r^2 = 0.99$.

been observed (Schaffer and Gaye, 1989; Larson et al., 1992; Pongsomboon et al., 1992). Hence, for avocado and mango trees grown in containers there is relative consistency between the reported values of Q_A and A_{max} .

The relatively low A_{max} rates that we observed for container-grown compared to field-grown trees may be attributed to containers restricting the root sink, thus causing the photoassimilate supply to exceed the capacity of demand (i.e., end-product inhibition of photosynthesis) (Schaffer et al., 1994). This is supported by

the increased concentrations of leaf starch for both crops growing in containers (160% to 230% higher in container-grown compared with NCS, field-grown trees). Increased leaf starch in avocado accompanied by a reduction in A_{max} was reported by Schaffer et al. (1987) who concluded that the latter occurred as a result of feedback inhibition. For other species, reduced *A* of container-grown plants as a result of root restriction has been attributed to end-product inhibition of photosynthesis caused by root restriction (Arp, 1991; Thomas and Strain, 1991). Although only starch was measured in this study, other nonstructural carbohydrates such as glucose and hexoses may have accumulated in the leaves as a result of sink (root) restriction. It has been recently proposed that accumulation of these end-products, particularly hexoses, in the leaves may repress genes that code for rubisco, resulting in feedback inhibition of *A* (Drake et al., 1997; Koch, 1996; Stitt, 1991). The hard, oolitic limestone soils of Florida can severely limit root growth, thus restricting the development of this sink (Crane et al., 1994; Schaffer et al., 1994). Thus, the relatively low A_{max} previously observed for field-grown trees in Florida may also be the result of end-product inhibition of photosynthesis.

The lower Q_A and A_{max} of field-grown avocado and mango trees that were determined for CS compared to NCS trees, were most likely due to photoinhibition of *A* as a result of low temperatures interacting with high incident PPF (Groom et al., 1991). Limited exposure of tropical and subtropical species to below 10 °C can result in reversible inhibition of photosynthesis (Taylor and Rowley, 1971). The F_v/F_m ratios for leaves of NCS, field-grown trees were 0.81 ± 0.02 and 0.80 ± 0.03 , for avocado and mango, respectively (Table 2), indicating that the photosynthetic processes were functioning normally in autumn (Öquist and Wass, 1988). The F_v/F_m ratios for avocado and mango trees were lower in winter when minimum daily temperatures decreased below 10 °C compared to early fall when chilling did not occur. Leaves also visibly appeared chlorotic following exposure to winter temperatures suggesting photooxidation of the chlorophyll. The lower F_v/F_m ratios indicated a reduced photochemical conversion efficiency of PS II (Krause, 1988), which can be an effect of photoinhibition at chilling temperatures (Groom et al., 1991). Photoinhibition from chilling injury can reduce Φ and Q_A (Powles et al., 1983) as we observed for avocado and mango during the winter measurement period. For both species, reduced *A* in winter may also have been partially a result of feedback inhibition. Since the canopy of avocado and mango trees is relatively quiescent during winter, the resultant reduction in sink strength promotes an increase in starch concentrations in leaves (Whiley, 1994). In our study, leaf starch

Table 2. The effect of growing conditions on CO₂ saturation of CO₂ assimilation (C_{aSAT}), the maximum rate of CO₂ assimilation at saturating partial pressures of CO₂ (A_{maxCa}), chlorophyll fluorescence (F_v/F_m) and leaf starch concentrations on a dry weight basis. Values of C_{aSAT} and A_{maxCa} were estimated from the regression models presented in Figs. 3 and 4 which were based on pooled data from one leaf of each of five trees. F_v/F_m and leaf starch data are mean values \pm standard errors (n = 5).

Treatment	C_{aSAT} ($\mu\text{mol}\cdot\text{mol}^{-1}$)	A_{maxCa} ($\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$)	F_v/F_m	Leaf starch concn ($\text{mg}\cdot\text{g}^{-1}$)
Avocado				
NCS field-grown	1473	50.6	0.81 ± 0.02	17.4 ± 0.14
CS field-grown	1357	34.9	0.41 ± 0.03	23.4 ± 0.20
Container-grown	1261	26.1	---	28.0 ± 0.26
Mango				
NCS field-grown	1113	30.3	0.80 ± 0.03	14.0 ± 0.03
CS field-grown	636	8.3	0.45 ± 0.03	25.4 ± 0.14
Container-grown	871	21.5	---	32.2 ± 0.18

concentrations in CS, field-grown avocado and mango trees were higher than in NCS, field-grown trees, thereby creating some doubt about relative importance of photoinhibition to the reduction of *A*. Further investigations to isolate the effect of cold stress from root restriction and elevated carbohydrate concentrations are required to clarify the situation.

The positive linear correlation between C_i and C_a implied that CO_2 diffusion through the mesophyll tissue was never limiting *A*. Leaves of NCS, field-grown trees had higher C_{aSAT} than leaves from CS, field-grown and container-grown trees. The carbon source–sink balance is a major factor in determining the response of plants to elevated CO_2 (Arp, 1991). For example, reduced *A* of bean plants in a CO_2 -enriched environment has been positively correlated with the carbon source–sink ratio (Peet, 1984). Thus, the lower C_{aSAT} for container-grown than for field-grown avocado and mango presumably was from an increased source–sink ratio due to root restriction. For field-grown trees in winter, either photoinhibition, feedback inhibition of *A* due to the lack of vegetative growth, or their combined effects may have resulted in a downward regulation of *A* and lower C_{aSAT} . However, C_{aSAT} can also be reduced through source-imposed limitations to *A* such as a reduction of incident PPF (Ehret and Joliffe, 1985).

When comparing the relative efficiency and capacity of *A* between the two species in this study, the response of avocado to varying photon and CO_2 fluxes was greater, 25% and 67%, respectively, than for mango. These differences may be attributed to the contrast in leaf longevity between the two species. For many species the photosynthetic rate is negatively correlated to life span of the leaf, with deciduous species in general having higher net photosynthetic rates than evergreen species (Chabot and Hicks, 1982; Larcher, 1969). At the evolutionary level a hypothesis advanced to explain this phenomenon is “longer leaf life spans compensate for environmentally limited photosynthetic activity” (Chabot and Hicks, 1982). Compared to mango trees, the physiology of avocado trees more closely resembles that of woody deciduous species. There is comparatively little carbon investment into avocado leaves that grow rapidly but have a longevity of <10 months. In contrast, mango leaves are very sclerophyllous with a high investment of carbon during growth and are sustained on trees for up to 5 years (Whiley and Schaffer, 1994). While avocado was shown to have a greater capacity to respond to increased C_a , this may not necessarily result in greater yield as there is a higher carbon investment in fruit growth of this oil-rich crop compared with sugar-accumulating species such as mango (Wolstenholme, 1986, 1987).

Literature Cited

Arp, W.J. 1991. Effects of source–sink relations on photosynthetic acclimation to elevated CO_2 . *Plant, Cell Environ.* 14:869–875.

Björkman, O. 1987. Low temperature chlorophyll fluorescence in leaves and its relationship to photon yield of photosynthesis photoinhibition, p. 123–144. In: D.J. Kyle, C.B. Osmond, and C.J. Arntzen (eds.). *Topics in photosynthesis*. vol. 9. Elsevier Press, Amsterdam.

Bower, J.P., B.N. Wolstenholme, and J.M. de Jagar. 1978. Incoming solar radiation and internal water status as stress factors in avocado, *Persea americana* Mill. cv. Edranol. *Crop Prod.* 7:129–133.

Chabot, B.F. and D.J. Hicks. 1982. The ecology of leaf life spans. *Annu. Rev. Ecol. Syst.* 13:229–259.

Chacko, E.K., P. Lu, and R.R. Kohli. 1995. Photoassimilate production and distribution in relation to productivity in mango (*Mangifera indica*), p. 109–124. In: *Mango 2000—Marketing seminar and production workshop proceedings*. Queensland Dept. of Primary Ind., Brisbane.

Crane, J.H., C. Balerdi, R. Campbell, and S. Goldweber. 1994. Managing fruit orchards to minimize hurricane damage. *HortTechnology* 4:21–27.

Demmig, B. and O. Björkman. 1987. Comparison of the effect of excessive light on chlorophyll fluorescence (77k) and photon yield of O_2 in leaves of higher plants. *Planta*

171:171–84.

Drake, B.G., M.A. Gonzales-Meler, and S.P. Long. 1997. More efficient plants: A consequence of rising atmospheric CO_2 . *Annu. Rev. Plant Physiol. Mol. Biol.* 48:609–639.

Ehleringer, J. and O. Björkman. 1977. Quantum yields for CO_2 uptake in C_3 and C_4 plants. *Plant Physiol.* 59:86–90.

Ehret, D.L. and P.A. Joliffe. 1985. Photosynthetic carbon dioxide exchange of bean plants grown at elevated carbon dioxide concentrations. *Can. J. Bot.* 63:2026–2030.

Groom, Q.J., N.R. Baker, and S.P. Long. 1991. Photoinhibition of holly (*Ilex aquifolium*) in the field during the winter. *Physiol. Plant.* 83:585–590.

Jones, H.G. 1992. *Plant and the Microclimate*. 2nd. ed. Cambridge Univ. Press, Cambridge.

Koch, K.E. 1996. Carbohydrate-modulated gene expression in plants. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 47:509–540.

Krause, G.G. 1988. Photoinhibition of photosynthesis. An evaluation of damaging and protective mechanisms. *Physiol. Plant.* 74:566–574.

Larson, K.D., B. Schaffer, and F.S. Davies. 1992. Flooding, mineral nutrition and gas exchange of mango trees. *Scientia Hort.* 52:113–124.

Larcher, W. 1969. The effect of environmental and physiological variables on the carbon dioxide gas exchange of trees. *Photosynthetica* 3:167–198.

Öquist, G. and R. Wass. 1988. A portable microprocessor operated instrument for measuring chlorophyll fluorescence kinetics in stress physiology. *Physiol. Plant.* 73:211–217.

Osman, A.M. and F.L. Milthorpe. 1971. Photosynthesis of wheat leaves in relation to leaf age, illuminance and nutrient supply. II. Results. *Photosynthetica* 5:61–68.

Peet, M.M. 1984. CO_2 enrichment of soybeans. Effects of leaf/pod ratio. *Physiol. Plant.* 60:38–42.

Pongsomboon, W., A.W. Whiley, R.A. Stephenson, and S. Subhabrabandhu. 1992. Effects of air temperatures on diurnal variation of water potential, conductance and CO_2 assimilation of mango (*Mangifera indica* L.) leaves. *Acta Hort.* 321:472–481.

Powles, S.B. 1984. Photoinhibition of photosynthesis induced by visible light. *Annu. Rev. Plant Physiol.* 35:15–44.

Powles, S.B., J.A. Berry, and O. Björkman. 1983. Interaction between light and chilling temperature on the inhibition of photosynthesis in chilling-sensitive plants. *Plant Cell Environ.* 6:117–23.

Rasmussen, T.S. and R.J. Henry. 1990. Starch determination in horticultural plant material by an enzymatic colorimetric procedure. *J. Sci. Food Agr.* 53:159–170.

Schaffer, B. and P.C. Andersen. 1994. Introduction, p. 1–2. In: B. Schaffer and P.C. Andersen (eds.). *Environmental physiology of fruit crops*. vol. 2. Sub-tropical and tropical crops. CRC Press, Inc., Boca Raton, Fla.

Schaffer, B. and G.O. Gaye. 1989. Net gas exchange and chlorophyll and nitrogen content of mango leaves as influenced by developmental light environment. *HortScience* 24:507–509.

Schaffer, B., L. Ramos, and S.P. Lara. 1987. Effect of fruit removal on net gas exchange of avocado leaves. *HortScience* 22:925–927.

Schaffer, B., C. Searle, A.W. Whiley, and R.J. Nissen. 1996. Effects of atmospheric CO_2 enrichment and root restriction on leaf gas exchange and growth of banana (*Musa*). *Physiol. Plant.* 97:685–693.

Schaffer, B., A.W. Whiley, and J.H. Crane. 1994. Mango, p. 165–211. In: B. Schaffer and P.C. Andersen (eds.). *Environmental physiology of fruit crops*. vol. 2. Sub-tropical and tropical crops. CRC Press, Inc., Boca Raton, Fla.

Schaffer, B., A.W. Whiley, and R.R. Kohli. 1991. Effects of leaf age on gas exchange characteristics of avocado (*Persea americana* Mill.). *Scientia Hort.* 48:21–28.

Scholefield, P.B., J.J. Walcott, P.E. Kriedeman, and A. Ramadanan. 1980. Some environmental effects on photosynthesis and water relations of avocado leaves. *Calif. Avocado Soc. Yrbk.* 64:93–105.

Searle, C., A.W. Whiley, D.R. Simpson, and J.B. Saranah. 1995. A preliminary phenophysiological model for ‘Kensington’ mango in subtropical Australia, p. 127–134. In: *Mango 2000—Marketing seminar and production workshop proceedings*. Queensland Dept. of Primary Ind., Brisbane.

Smillie, R.M., S.E. Hetherington, J. He, and R. Nott. 1988. Photoinhibition at chilling temperatures. *Austral. J. Plant Physiol.* 15:207–222.

Stitt, M. 1991. Rising CO_2 levels and their potential significance for carbon flow in photosynthetic cells. *Plant Cell Environ.* 14:741–762.

Syvrtsen, J.P. 1984. Light acclimation in citrus leaves. II. CO_2 assimilation and light, water, and nitrogen use efficiency. *J. Amer. Soc. Hort. Sci.* 109:812–817.

Taylor, A.O. and J.A. Rowley. 1971. Plants under climatic stress. I. Low temperature, high-light effects on photosynthesis. *Plant Physiol.* 47:713–718.

Thomas, R.B. and B.R. Strain. 1991. Root restriction as a factor in photosynthetic acclimation of cotton seedlings grown in elevated carbon dioxide. *Plant Physiol.* 96:627–634.

Whiley, A.W. 1984. Crop management. Review paper, p. 186–195. In: *Proc. 1st Austral. Mango Res. Wkshp.*, Cairns.

Whiley, A.W. 1994. Ecophysiological studies and tree manipulation for maximization of yield potential in avocado (*Persea americana* Mill.). PhD thesis, Univ. Natal, Pietermaritzburg, S. Afr.

Whiley, A.W. and B. Schaffer. 1994. Avocado, p. 3–35. In: B. Schaffer and P.C. Andersen (eds.). *Environmental physiology of fruit crops*. vol. 2. Sub-tropical and tropical crops. CRC Press Inc., Boca Raton, Fla.

Whiley, A.W., J.B. Saranah, B.W. Cull, and K.G. Pegg. 1988. Manage avocado tree growth cycles for productivity gains. *Queensland Agr. J.* 114:29–36.

Wolstenholme, B.N. 1986. Energy costs of fruiting as a yield-limiting factor with special reference to avocado. *Acta Hort.* 175:121–126.

Wolstenholme, B.N. 1987. Theoretical and applied aspects of avocado yield as affected by energy budgets and carbon partitioning. *S. Afr. Avocado Growers’ Assn. Yrbk.* 10:58–61.