

Managing *Phytophthora cinnamomi*


Liz Dann, Ken Pegg, Luke Smith
QPIF, Indooroopilly, QLD

Tony Whiley
Sunshine Horticultural Services



Topics covered

- Experimental results
 - Rootstock trials
 - Optimisation of phosphonate applications
- Integrated control



Phytophthora root rot (PRR)




PRR linked to seasonal physiology of the tree

- temperature & soil saturation play a major role in PRR severity

- Optimum temperature for
 - PRR development 19 – 25°C
 - Avocado growth 21 – 33°C
- < 22 °C root growth slows = PRR severe
- > 27 °C avocado grows better than P_c = PRR less severe
- P_c stress on trees greatest in spring

Effect of rootstock on PRR



Velvick clone (7.7)
Reed (8.3)


SHSR-04 (2.7)



Clonal Hass (4.8)




Duranbah rootstock trial, tree health



This work is continuing...



Optimising phosphonates



How do phosphonates work?

Dual action:

- High root levels – direct antimicrobial action
- Low root levels – plant defence responses activated
- Moves with photosynthates, accumulates and persists in plant tissue with greatest metabolic activity at time of application

Methods of phosphonate application

- Trunk injections
- Foliar sprays
- Bark application
- Soil application

Phosphonate applications must be continued for the life of the tree

Trunk injection

- Most effective treatment – 20% PO_3 injected at even spacing around the trunk (15mL per metre canopy diam.)
- Central wood retains its transport function (unique in avocados)
- Injury to xylem near injection site
 - Becomes non-conductive
 - Replaced by tree in 1 – 2 years
- Healthy trees – 1 annual injection at maturation of summer flush
- Sick trees – 2 injections - at maturation of both summer and spring flushes

Foliar Sprays


- Achieve same root concentrations as injections
- Need to monitor root levels to determine number of applications necessary
- 0.5% (8.3mL/L of the 600 product)
- More sprays required to achieve 0-50ppm, than >50ppm
- No applications within 6 weeks of flowering

Bark Sprays

- Highly effective in young trees where stem has some photosynthetic (green) activity
- Very effective against trunk cankers
- Unable to achieve and maintain adequate root levels in older trees, unless applied frequently
- 20% product with 2% bark penetrant
 - Higher than 20% causes chemical cincturing


Soil Drenching

- Initial increase in root concentration but quickly declines
- Need to apply monthly to maintain root levels
- Moves in xylem with water transport, thus not effectively distributed to target tissues




Phosphonate - inject vs trunk spray

Root levels



Later maturing varieties eg. Reed



Inject, bark, foliar

Foliar

Think about using metalaxyl

- Upward movement in transpiration stream – taken up only by roots
- Prevents colonisation of roots by zoospores (major)
- Toxic to germinating chlamydospores (minor)
- If feeder roots healthy, will be absorbed before degradation by soil micro-organisms
- Off patent (cheaper)



Integrated control

- Drainage
- Nursery practices
- Cultural control
- Resistance
- Chemicals
 - Phosphonates
 - Metalaxyl



Acknowledgements

**Thank you to the many growers
and collaborators for supporting
our research!**



Know-how for Horticulture™

