

The Effect of Indole-3-acetic Acid and Other Growth Regulators on the Ripening of Avocado Fruits¹

Received for publication June 17, 1974 and in revised form January 31, 1975

PETER O. TINGWA² AND ROY E. YOUNG

Department of Plant Sciences, University of California, Riverside, California 92502

ABSTRACT

Observations were made of the effects of several plant regulators, indole-3-acetic acid, kinetin, abscisic acid, and gibberellic acid, as well as of extracts prepared from leaves and fruit stalks on the respiration pattern, ethylene production, and the number of days to ripen of avocado fruits (*Persea americana* Mill.). These substances were vacuum infiltrated to insure good penetration and distribution. Kinetin, abscisic acid, gibberellic acid, and the extracts had no effect on either ripening time or on the respiration pattern and ethylene production of the fruits. Indoleacetic acid, however, had a marked effect on ripening. At high concentrations (100 and 1000 μM), indoleacetic acid stimulated respiration and induced preclimacteric ethylene production, resulting in accelerated ripening of the fruits. At the low concentrations (1 and 10 μM), it delayed ripening of fruits and suppressed the climacteric respiration and ethylene production. The results reinforce several previous observations with other fruits that auxins may largely constitute 'resistance to ripening' and may be responsible for the lack of ripening shown by unpicked fruits.

The inability of some fruits, notably the avocado, to ripen on the tree (3, 7, 8), coupled with the lack of response by freshly harvested fruits to ethylene treatment (16, 20, 23), have prompted several workers to propose the existence of a ripening inhibitor in attached fruits (3, 7, 8). This inhibitor is presumed to be formed in leaves of the parent tree and translocated to the fruit while still attached (3, 7, 8). Following harvest, the inhibitor is inactivated during the preclimacteric period and ripening is initiated by endogenously produced ethylene (14, 16, 20, 23). Plant hormones have been shown to affect some aspects of fruit ripening. For example, indoleacetic acid has been reported to delay the onset of the climacteric in bananas (30) and pears (14), and both gibberellins and cytokinins have been shown to delay the degradation of Chl in ripening tomatoes (1) and orange peel (12).

A growing body of information indicates that the ripening inhibitor may be an auxin (14, 15, 27). Direct evidence that auxins constitute the resistance to ripening was demonstrated by the delay of ripening processes in bananas (30) and pears (14) following treatments with either 2,4-dichlorophenoxy-

acetic acid or indoleacetic acid. IAA-oxidase activity has been demonstrated in ripening fruit (13, 14, 15) and is presumed to play a physiological role in the regulation of auxin level.

Because attached avocado fruit resist ripening for long periods (3, 7, 8), we decided to test the effects of hormones and regulators as well as extracts of leaves and stems on the ripening of detached avocado fruits.

MATERIALS AND METHODS

Source of Fruits. Avocado fruits (*Persea americana* Mill.) of both 'Fuerte' and 'Hass' cultivars were obtained from the South Coast Field Station of the University of California. For each experiment conducted, uniform fruits were picked from a single tree to minimize variability.

Infiltration of Fruits. Infiltration was carried out according to the procedure of Frenkel and Dyck (14) with slight modifications. Pressure over the fruits was reduced to 260 mm Hg for 6 min and infiltration time was reduced to 5 min to minimize the development of anaerobic conditions. When intact fruits were used, the peduncle was removed to improve penetration of the infiltrated substances. Infiltrated fruit was compared with water-infiltrated as well as uninfiltrated fruit. In some experiments, fruits were cut in half; one half being infiltrated with a solution of the test substance and the other with water. On the average, an increase in weight of about 3% was obtained from infiltration.

Determination of CO₂ and Ethylene. Fruits were placed singly in 1.8-liter jars at 20 C. Humidified air was passed at 70 ml/min and respiration followed with a Beckman infrared gas analyzer (Model 215A). Ethylene production was monitored automatically every 3 hr on an Aerograph HY-FI Model 600-D gas chromatograph using a 180 × 0.3 cm column of Porapak Q. Output of the chromatograph was printed by a Varian Model 480 digital integrator.

Growth Regulator Treatments. Fruits were treated with solutions of IAA, GA₃, ABA, kinetin, and extracts of leaves and peduncles. Extracts were made by blending 20 g fresh weight in 200 ml of 80% ethanol. The homogenate was filtered and the volume of the filtrate reduced to 20 ml under reduced pressure at 40 C. The concentrate was centrifuged at 4500g and stored at -20 C. Treatment solutions were made by diluting the concentrate with 4 or 8 volumes of redistilled H₂O.

RESULTS

Vacuum Infiltration. Subjecting the fruit to vacuum treatment with or without infiltration of water or aqueous solution caused an immediate and marked reduction of respiration for 2 to 3 hr, followed by an increase in respiration of 3 to 4 times normal which persisted for 1 to 2 hr. (These short term changes in respiration rate are not shown in either Fig. 1a or

¹ This study represents part of the Ph.D. dissertation submitted by P.O.T.

² Present address: Faculty of Agriculture, P. O. Box 32, Khartoum, North, Sudan.

2a). The cause of the burst of CO_2 production is presumed to be attributable to either an injury effect, or a sudden increased availability of O_2 to an accumulated oxidizable substrate. Although the practice of vacuum infiltration enhanced preclimacteric respiration, it caused the opposite effect for climacteric respiration.

IAA Effect. The influence of increasing concentration of IAA from 1 to 1000 μM on the respiration pattern of 'Fuerte' and 'Hass' fruits is shown in Figures 1a and 2a, respectively. When compared to water-infiltrated controls, preclimacteric respiration was stimulated by IAA and the magnitude of stimulation was proportional to its concentration. The reduced respiration rate at the climacteric resulting from the practice of infiltration was reversed by higher IAA concentrations (100 and 1000 μM), such that the climacteric respiration in these treatments was as high as that of uninfiltrated controls. On the other hand, low IAA concentrations (1 and 10 μM), while stimulating preclimacteric respiration, depressed the climacteric respiration rate even further.

The effect of IAA on ethylene production is shown in Figures 1b and 2b for 'Fuerte' and 'Hass' varieties, respectively. High concentrations of IAA induced preclimacteric ethylene production earlier and stimulated its rate of evolution during the climacteric. The peak production of ethylene by fruits given these higher IAA treatments came before those of the controls. Low IAA levels (1 and 10 μM) caused a delay in initiation of ethylene production during the preclimacteric period and both delayed and reduced its production at the climacteric peak.




FIG. 1. Effect of increasing concentration of IAA on respiratory pattern (a) and ethylene production (b) of 'Fuerte' avocado fruits at 20 C. The IAA concentrations were 1 μM (Δ), 10 μM (\blacktriangle), 100 μM (\square), 1000 μM (\blacksquare), and uninfiltrated (\circ) and water-infiltrated (\bullet) controls. Each point represents the average of four fruits.




FIG. 2. Respiration pattern (a) and ethylene production (b) of 'Hass' avocado fruits vacuum infiltrated with water (\bullet), 1 μM (Δ), 5 μM (\blacksquare), and 10 μM (\blacktriangle) IAA at 20 C. Uninfiltrated fruits are presented as (\circ). Each point is the average of four fruits.

When judging the effect of IAA (or any of the other treatments) on the time of ripening, the days preceding the climacteric ethylene peak were considered to be the number of days taken to ripen because this point is related to ripening and may be determined with great accuracy. Therefore, as shown in Figures 1b and 2b, low concentrations of IAA delayed ripening of fruits by as much as 2 to 3 days at 20 C whereas high concentrations accelerated ripening. This delay of ripening induced by low IAA concentrations was much more pronounced in the midseason 'Fuerte' (Fig. 1b) than in late season 'Hass' fruits (Fig. 2b).

Kinetin, GA₃, ABA, and Extracts. The respiration pattern and ethylene production of fruits infiltrated with 1 or 10 μM solutions of GA₃, kinetin, ABA, or with 1:4 and 1:8 dilutions of leaf or peduncle extracts did not differ from uninfiltrated or water-infiltrated controls. There was, however, some preclimacteric stimulation of ethylene production by fruits infiltrated with 10 μM kinetin.

The effects of these treatments on ripening time are given in Table I. Although there was some variability in the ripening time of individual fruits, none of the concentrations of these treatments affected the time of ripening. In contrast, the ripening time of fruits infiltrated with 1 μM IAA was uniform and delayed by 2.5 days when compared to the controls.

Results of further investigations with half fruits showed no significant difference between ripening time of halves treated with 10 μM ABA or 1:4 peduncle extract and their corresponding water-treated halves. In each case, the respiration pattern and ethylene production were similar in both halves of the

fruit as shown in Figure 3, a and c. In contrast, whenever one half of a fruit was treated with $1 \mu\text{M}$ IAA, its respiration and ethylene production were suppressed and its ripening was delayed (Fig. 3b).

DISCUSSION

These experiments show that IAA exerts a marked effect on the ripening of detached avocado fruits, thus providing the first indications of the involvement of auxins (IAA) in the ripening of avocado fruits. They support previous reports for pears (14) and banana slices (30) that auxins can delay the ripening of fruits, provided adequate penetration of the substance is insured by infiltration. Opposite effects, resulting from poor penetration of auxins, have been reported (5, 19) and alluded to (14, 30) in pears and bananas.

Treatment of avocado fruit with IAA seems to exert control over the time of induction of the climacteric as well as over the amount of ethylene produced during the climacteric.

Table I. Effect of Various Hormones and Leaf and Peduncle Extracts on the Ripening Time of 'Hass' Avocado Fruits at 20 C

The number of days shown are averages of four fruits.

Substance Infiltrated	Days to Ripen
None	6.8
Water	6.7
$1 \mu\text{M}$ IAA	9.1
$1 \mu\text{M}$ Kinetin	7.0
$10 \mu\text{M}$ Kinetin	6.8
$1 \mu\text{M}$ GA ₃	7.0
$10 \mu\text{M}$ GA ₃	6.8
$1 \mu\text{M}$ ABA	6.8
$10 \mu\text{M}$ ABA	6.6
1:8 Leaf extract	7.1
1:4 Leaf extract	6.7
1:8 Peduncle extract	6.9
1:4 Peduncle extract	7.4

Concentrations of IAA above $10 \mu\text{M}$ caused more rapid ripening and higher ethylene production, an effect reported for auxin treatments in other fruits as well (9, 14, 19, 30). IAA concentrations less than $10 \mu\text{M}$ caused delayed ripening and ethylene production lower than the controls. These seemingly contradictory observations suggest that IAA is acting on two different systems. Dual effects of auxin have been proposed in other cases, e.g., pea root tips (28) in which IAA affected two systems influencing ethylene synthesis; the low concentration treatment was shown to act directly on an enzyme whereas the high concentration system required protein synthesis. Likewise in leaf abscission experiments, IAA plays a dual role in that, in general, auxin promotes abscission at low treatment levels and inhibits at higher levels (2). In the case of IAA effects on stem elongation, stimulation at low and inhibition at high levels is the rule (29). Vendrell (30) has suggested that auxins affect two regulatory systems in bananas; one concerned with regulation of respiration and ethylene production and the other with desensitizing fruits to the effects of ethylene.

We propose a dual effect for IAA in avocado ripening. The high concentration effect is to induce early ethylene production which in turn initiates ripening. It may be noted in Figure 1 that the first increase in ethylene production precedes the increase in respiration by a few hours when $1000 \mu\text{M}$ IAA was used. On the other hand, the first increase in ethylene production in the $1 \mu\text{M}$ -treated fruit and in the control fruit occurred about 12 hr after respiration increased. We suggest that treatment with $1000 \mu\text{M}$ IAA caused increased and earlier ethylene production, which initiated ripening just as does treatment with ethylene after picking (4, 16). Ethylene production in response to auxin treatment has been reported for a number of plant tissues (24).

The low concentration effect of IAA is clearly at a different site because ethylene production was both delayed and diminished, relative to the control (Fig. 1). Assuming that the low IAA concentration effect mimics the process by which avocados remain unripe on the tree, different mechanisms may be visualized for the effect of IAA, depending on which of two alternatives are assumed: (a) a ripening inhibitor is translocated from tree to fruit; or (b) a ripening hormone produced in the fruit is kept at a low level by movement into the




FIG. 3. Respiration pattern and ethylene production of 'Fuerte' half fruits infiltrated with (a): $10 \mu\text{M}$ ABA (●), (b): $1 \mu\text{M}$ IAA (■), and (c): 1:4 peduncle extract (▲), all at 20 C. The other halves of the same fruits infiltrated with water are shown as (○).

branch before detachment and ripening is initiated by accumulation of the hormone after detachment. If ripening in attached fruit is prevented by the first alternative and ripening is initiated by depletion of IAA, treatment of the detached fruit with low levels of IAA would contribute to the endogenous pool and extend the preclimacteric period. This is so as long as the IAA concentration is below the level that induces ethylene production. If the second alternative is the case, *i.e.*, ripening initiated by a hormone accumulating in detached fruit, low concentrations of IAA must delay ripening by competing with the hormone for active sites or by acting as a negative effector on the ripening control mechanism.

Neither kinetin nor GA had any effect on the time of ripening of avocado fruits, as shown in Table 1. They also had no effect on the respiration pattern or ethylene production of fruits except for noticeable stimulation of ethylene production by 10 μ M concentration of kinetin. The roles of these two regulators in delaying the ripening of some fruits is well documented (1, 11, 12, 25, 31). Gibberellins have been shown to delay ripening of bananas (25, 31) and tomatoes (1, 11). Because gibberellins and ethylene have opposing effects (26), this group of hormones has been considered, together with auxins, to control the ripening of fruits (27).

The effects of gibberellins and cytokinins may not delay all aspects of normal ripening but affect only one or a few criteria of ripening. In several cases, only pigment changes of chlorophyll breakdown, or lycopene synthesis have been affected by these hormones (1, 11, 22). Blumenfeld and Gazit (6) and Gazit and Blumenfeld (17) have demonstrated the existence of gibberellins and cytokinins, respectively, in avocado fruits. They have also pointed to the possible synthesis of these hormones in the fruit, particularly in the seed coat. The lack of response of the avocado fruit to exogenous kinetin and GA₃ may be attributable to adequate endogenous amounts of these hormones. In addition, because cell division continues throughout the life of the avocado fruit, a nonlimiting amount of these hormones, especially cytokinins, in detached fruit is possible.

ABA did not seem to influence the ripening time of intact 'Hass' fruits (Table I) nor respiration pattern or ethylene production of 'Fuerte' half fruits (Fig. 3a). This was unexpected because ABA is considered to be a senescence-promoting hormone and was thus assumed to hasten the ripening of the fruits. In tomatoes, ABA accelerated ripening (21, 22) by increasing respiration rate and ethylene production (21), as well as by stimulating lycopene synthesis (22). Indirect evidence linking ABA to fruit ripening has also been provided by Goldschmidt *et al.* (18) who reported that an increase in ABA-like inhibitors accompanied senescence of citrus fruits. Information on the role of ABA in ripening of horticultural fruits is still meager; as a result, the extent of its participation in fruit ripening is not yet clear (10).

The results in Table 1 and Figure 3c show that extracts prepared from leaves and peduncle had no effect on the ripening of the fruit. This failure of the extracts to elicit any effect does not necessarily indicate the absence of the ripening inhibitor but rather may be attributable to a weakness of the extraction procedure adopted.

We conclude that the extent of the influence of IAA on avocado ripening by far exceeds that of the other hormones studied. Auxins are likely to be the substances (inhibitors) that prevent ripening of unpicked or freshly harvested avocado fruits. The significance of optimum balance between auxins and the other hormones for the effect to be shown cannot be discounted. The unequivocal identification of this substance

must come not only from extraction from the plant but also from confirmation of its effect on unpicked fruits.

Acknowledgment—We thank Drs. J. B. Biale and L. C. Erickson for helpful suggestions and for reading the manuscript. The technical assistance of Cameron Duncan is greatly appreciated.

LITERATURE CITED

1. ABDEL KADER, A. S., L. L. MORRIS, AND E. C. MAXIE. 1966. Effect of growth regulating substances on the ripening and shelf-life of tomatoes. *Hortsci.* 1: 90.
2. ADDICOTT, F. T. 1970. Plant hormones in the control of abscission. *Biol. Rev.* 45: 485-524.
3. BIALE, J. B. 1960. Respiration of fruits. *Handbuch der Pflanzen-Physiologie.* 12: 536-592.
4. BIALE, J. B. AND R. E. YOUNG. 1971. The avocado pear. In: A. C. Hulme, ed., *The Biochemistry of Fruits and their Products*. Academic Press, New York, p. 1-63.
5. BLAKE, J. R. AND C. D. STEVENS. 1959. Effects of growth regulating substances on the ripening of the banana fruits. *Queensl. J. Agric. Sci.* 16: 87-90.
6. BLUMENFELD, A. AND S. GAZIT. 1972. Gibberellin-like activity in developing avocado. *Physiol. Plant.* 27: 116-121.
7. BURG, S. P. 1964. Studies on the formation and function of ethylene gas in plant tissue. In: J. P. Nitsch, ed., *Régulateurs Naturels de la Croissance Végétale*. Centre National de la Recherche Scientifique, Paris. pp. 719-725.
8. BURG, S. P. AND E. A. BURG. 1964. Evidence of a naturally occurring inhibitor of ripening. *Plant Physiol.* 39: S-x.
9. BURG, S. P. AND E. A. BURG. 1965. Relationship between ethylene production and ripening in bananas. *Bot. Gaz.* 129: 200-204.
10. DILLEY, D. R. 1969. Hormonal control of fruit ripening. *Hortsci.* 4 (2): 111-113.
11. DOSTAL, H. C. AND A. C. LEOPOLD. 1967. Gibberellin delays ripening of tomatoes. *Science* 158: 1579-1580.
12. EILATI, S. K., E. E. GOLDSCHMIDT, AND S. P. MONSELISE. 1969. Hormonal control of colour changes in orange peel. *Experientia* 25: 209.
13. FRENKEL, C. 1972. Involvement of peroxidase and indole-3-acetic acid oxidase isozymes from pear, tomato, and blueberry fruit ripening. *Plant Physiol.* 49: 757-763.
14. FRENKEL, C. AND R. DYCK. 1973. Auxin inhibition of ripening in Bartlett pears. *Plant Physiol.* 51: 6-9.
15. FRENKEL, C. AND N. F. HARD. 1973. Initiation of ripening in Bartlett pear with an antiauxin α (*p*-chlorophenoxy)isobutyric acid. *Plant Physiol.* 52: 380-384.
16. GAZIT, S. AND A. BLUMENFELD. 1970. Response of mature avocado fruits to ethylene treatment before and after harvest. *J. Amer. Soc. Hort. Sci.* 95: 229-231.
17. GAZIT, S. AND A. BLUMENFELD. 1970. Cytokinin and inhibitor activities in avocado fruit mesocarp. *Plant Physiol.* 46: 334-336.
18. GOLDSCHMIDT, E. E., S. K. EILATI, AND R. GOREN. 1970. Increase in ABA-like inhibitors and decrease in GA-like substances during ripening and senescence of citrus fruits. In: J. D. Carr, ed., *Plant Growth Substances*. Springer-Verlag, Berlin. pp. 611-617.
19. HANSEN, E. 1946. Effect of 2,4-dichlorophenoxyacetic acid on the ripening of Bartlett pears. *Plant Physiol.* 21: 588-592.
20. HANSEN, E. 1966. Postharvest physiology of fruits. *Annu. Rev. Plant Physiol.* 17: 459-480.
21. KADER, A. A., L. L. MORRIS AND J. M. LYONS. 1973. Effect of abscisic acid on ripening of tomato fruits. *Hortsci.* 8: 264.
22. KHUDAIRI, A. K. 1972. The ripening of tomatoes. *Amer. Sci.* 60: 696-707.
23. MAPSON, L. W. 1970. Biosynthesis of ethylene and the ripening of fruit. *Endeavour* 29: 29-33.
24. PRATT, H. K. AND J. D. GOESCHL. 1969. Physiological roles of ethylene in plants. *Annu. Rev. Plant Physiol.* 20: 541-584.
25. RUSSO, L., JR., H. C. DOSTAL, AND A. C. LEOPOLD. 1968. Chemical regulation of ripening in bananas. *Bioscience* 18: 109.
26. SCOTT, P. C. AND A. C. LEOPOLD. 1967. Opposing effects of gibberellin and ethylene. *Plant Physiol.* 42: 1021-1022.
27. SMOCK, R. M. 1972. Influence of detachment from tree on the respiration of apples. *J. Amer. Soc. Hort. Sci.* 94: 509-511.
28. STEEN, D. A. AND A. V. CHADWICK. 1973. Effects of cycloheximide on indoleacetic acid induced ethylene production in pea root tips. *Plant Physiol.* 52: 171-173.
29. THIMANN, K. V. 1937. On the nature of inhibitions caused by auxin. *Amer. J. Bot.* 24: 407-412.
30. VENDRELL, M. 1969. Reversion of senescence: effect of 2,4-dichlorophenoxyacetic acid and indoleacetic acid on respiration, ethylene production and ripening of banana fruit slices. *Aust. J. Biol. Sci.* 22: 601-610.
31. VENDRELL, M. 1970. Acceleration and delay of ripening of banana fruit by gibberellin acid. *Aust. J. Biol. Sci.* 23: 553-559.