

AVOCADO PLANTING SYSTEMS

B. Faber¹, G. Thorp² and A. Barnett²

¹**University of California Cooperative Extension, Ventura Co, 669 County Square Dr.,
Ventura, CA, USA. E-mail: bafaber@ucdavis.edu**

²**HortResearch, Mt. Albert Research Centre, 120 Mt. Albert Rd., Private Bag 92 169,
Auckland, NZ. E-mail: gthorp@hort.cri.nz**

ABSTRACT

In many countries around the world recommendations are made for planting large, one meter by one meter by one meter holes that are backfilled with a mixture of native soil and 25% by volume of organic planting material. Aside from being expensive, this is very labor intensive. Trials were established in New Zealand on a well drained sandy loam, *Phytophthora*-free soil with seedling rootstocks and in California on a clay loam, infested with root rot using clonal rootstocks. 'Hass' scions were used. Trees were planted into large holes (1.5 X 0.8 X 0.6 m) with or without 25% well rotted compost planting mix or small holes (0.3 m wide and 0.45 m deep) with or without the organic planting mix. Trees were measured for stem girth, tree height and canopy dimension. After 20 months of growth in New Zealand and after 18 months growth in California there were no differences in tree dimensions with any of the combinations of hole size or backfill composition.

Key Words: *Phytophthora*, planting hole, soil amendment

INTRODUCTION

In numerous publications world-wide, planting hole recommendations for avocado and other sub-tropical crops are made for large holes from 0.6 by 0.6 by 0.6 meters to 1 cubic meter. These recommendations also include incorporation of manures or composts comprising 25% by volume with the native soil (Donadio, 1987; Teixeira *et al.*, 1992; Anon, 2003; Solares, 1981; Anon, 1982; Choo, 2002; Galan Suco, 1987). The authors have noted the use of large holes and amendments in several countries, including New Zealand, Guatemala, Brazil, Costa Rica, Mexico and the United States.

The various reasons given for making these large holes are to disrupt any compaction or limiting soil layers and to create a more conducive environment for root growth. In the case of replanting deciduous orchards, it has been found to be beneficial in actually replacing the native soil in the hole with pathogen free soil (McKenry, 1996). In many cases, research has shown that holes much larger than the planting ball and using organic amendments can cause problems for many tree species (Whitcomb, 1979; Harris *et al*, 2004). Improper mixing of the organic amendment can cause anaerobic conditions and settling due to amendment decomposition. Soil that has not been properly firmed in the hole can also lead to plant settling and stems can drop below grade leading to crown rot (Harris *et al*, 2004).

Nonetheless, on the basis of recommendations made in many countries there could be some value in these planting practices, especially in the light of the effect organic matter has on avocado root rot. Numerous studies have shown organic matter suppresses the causal agent of root rot (Downer, 2001). This study evaluates the effect of hole size and amendments on avocado growth in an ideal environment with excellent soil conditions and in a more harsh one with heavy soil texture and the presence of the root rot pathogen.

MATERIALS AND METHODS

On the north island of New Zealand at two sites in the Bay of Plenty, 20 trees each were planted to one of four treatments: a) small holes (30 x 45 cm) without amendment; b) small holes with 25% by volume compost; c) big holes (150 x 80 x 60 cm) without amendment and d) big holes with 25% by volume compost. Big holes were dug with a backhoe, while small holes were dug by shovel. Trees were approximately 0,65 m tall at planting. Soil was a deep sandy loam at both sites. Trees were irrigated by drip irrigation. Trees were 'Hass' on 'Zutano' seedling rootstock. Trees were planted the second week of spring 2000. Tree height, trunk caliper and canopy volume were measured on a monthly basis for eight months and then twice a year for the next year. In Carpinteria, California a similar trial was established using 'Hass' on 'Toro Canyon' rootstock. Trees were approximately 0.75 m tall at planting. The grove has a heavy clay loam soil and a history of root rot. The trees are on drip irrigation. The trees were planted summer 2001 and monitored for 18 months after planting.

RESULTS AND DISCUSSION

Figures 1-3 show the results of the different planting treatments at three sites, two in New Zealand on ideal soils and one on the heavy soil infected with root rot in California. Only tree height is shown; trunk girth and canopy volume followed similar patterns. From planting onwards, there were no differences in tree growth in any of the treatments at any of the sites. This would lead one to the conclusion that there is no value in and great expense in making big holes and incorporating amendment. This is especially so in hillside situations where moving equipment and amendments on steep slopes would be very difficult.

The trees at the Carpinteria site, although infested with root rot, all looked good. The addition of organic matter in conjunction with the clonal rootstocks did not apparently provide any greater disease resistance. This is in accordance with work done by John Menge (1997) which shows that the greatest benefit derived from mulching are seedling rootstocks. The effect of mulch on disease suppression diminishes with the rootstocks resistance to root rot.

REFERENCES

- ANONYMOUS. 1982. El cultivo del aguacate. 2nd ed. Federacion Nacional de Cafeteros de Colombia, Programa de Desarrollo y Diversificacion de Zonas Cafeteras. p.23.
- ANONYMOUS. 2003. National Dept of Ag in South Africa <http://www.nda.agric.za>
- CHOO, W.K. 2000. Longan production in Asia. FAO RAP Pub: 2000/20. Rome, Italy. p.126
- DONADIO, L.C. 1987. Present status of Brazilian avocado industry. Proceed. First World Avocado Cong. S. African Avocado Growers' Association Yearbook 1987. 10:82-85.
- DOWNER, A.J., J. A. MENGE, AND E. POND. 2001. Effects of cellulytic enzymes on *Phytophthora cinnamomi*. Phytopathology 91:839-846.
- HARRIS, R.W., J.R. CLARK AND N.P. MATHENY. 2004. Arboriculture. Prentice Hall. Upper Saddle River, New Jersey. p.578.
- GALAN SAUCO, V. AND U.G. MENINI. 1987. Litchi cultivation. FAO PPP Paper: 83. Rome, Italy. p.94.
- GONCALVES TEIXEIRA, C., E. W. BLEINROTH, J. VIDIGAL DE CASTRO, Z.J. DE MARTIN, J.S. TANGO, J. MENEGALDO TURATTI, R.S.S. FERNANDES LEITE, AND A.E. BRITO GARCIA. 1992. Abacate: cultura, material-prima, processamento e aspectos economicos. 2nd Ed. ITAL-Campinas. p. 250.
- MCKENRY, M.V. AND T. BUZO. GROWTH BENEFIT OF ADDING "VIRGIN SOIL" WHEN REPLANTING AN ORCHARD. PROCEEDINGS, AG FRESNO, NOVEMBER 1996. Proceedings, Ag Fresno, November 1996.
- MENGE, J.A. 1997. Biocontrol of Phytophthora root rot of avocado with mulch and biocontrol agents. Proceed. Calif Avo Research Symposium. Riverside, CA.
- RODRIGUEZ SUPPO, F. 1992. El Aguacate.. Edf. AGT, Mexico. p.167
- SOLARES, M. 1981. Tecnicas y practicas en el cultivo del aguacate.. 3rd ed. Mexico: Editores Mexicanos Unidos. p. 226.
- WHITCOMB, C.E. 1979. Amendments and tree establishment. J. Arboriculture. 5(10): 217-19.

Figure 1. Tree height at site 1 in New Zealand 20 months after planting. No differences were found at the 5% level of significance.




Figure 2. Tree height at site 2 in New Zealand 20 months after planting. No differences were found at the 5% level of significance.




Figure 3. Tree height in California 18 months after planting. No differences were found at the 5% level of significance.

