

## **COMMERCIAL RIPENING TRIALS WITH SOUTH AFRICAN AVOCADOS IN THE UNITED KINGDOM**

F.Kruger<sup>1</sup> and D. Lemmer<sup>1</sup>

<sup>1</sup>ARC-Institute for Tropical and Subtropical Crops, Private Bag X11208, Nelspruit 1200  
South Africa, Tel: +27 (0)13 753 7000, Fax: +27 (0)13 752 3854 Email: [fransk@arc.agric.za](mailto:fransk@arc.agric.za)

During July 2006, one container each of controlled atmosphere (CA) and SmartFresh<sup>SM</sup> Hass and Fuerte avocados were exported from Westfalia Estates to the United Kingdom. Each consignment contained 7 Hass and 3 Fuerte samples. The ripening patterns of the fruit were recorded by two prepackers. The controlled atmosphere fruit took an average of 4.98 days to ripen while the SmartFresh<sup>SM</sup> fruit required 5.44 days. Although considerable variation occurred between samples, certain of the cultivar/count combinations displayed very similar ripening patterns whether treated with SmartFresh<sup>SM</sup> or stored under CA. In both cultivars, the largest SmartFresh<sup>SM</sup> fruit ripened faster than the CA fruit while the reverse was true for the smallest counts. The results further confirmed our previous recommendation regarding the 70% moisture content (30% dry mass) cut off point for SmartFresh<sup>SM</sup> fruit destined for the 'Ripe and Ready' market sector. It may, however, be possible to send larger counts (>300g) before this point is reached while it may be advisable not to send very small counts (<150g) to ripeners. Although the incidence of greypulp (diffuse mesocarp discolouration) was low in the present trial, a commercial trial conducted during the previous season has clearly demonstrated the greypulp reducing capacity of SmartFresh<sup>SM</sup> (1-methylcyclopropene).

## **ENSAYOS COMERCIALES DE MADURACIÓN DE AGUACATES SUDAFRICANOS EN EL REINO UNIDO**

F.Kruger<sup>1</sup> y D. Lemmer<sup>1</sup>

<sup>1</sup>ARC-Institute for Tropical and Subtropical Crops, Private Bag X11208, Nelspruit 1200  
South Africa, Tel: +27 (0)13 753 7000, Fax: +27 (0)13 752 3854 Email: [fransk@arc.agric.za](mailto:fransk@arc.agric.za)

En julio de 2006, se exportaron en contenedores aguacates Hass y Fuerte, tratados con SmartFresh<sup>SM</sup> o en atmósfera controlada (AC), desde Westfalia Estates al Reino Unido. Cada envío contenía 7 muestras de la variedad Hass y 3 muestras de la variedad Fuerte. Los parámetros de maduración de la fruta fueron registrados por dos empaquetadores preliminares. La fruta en atmósfera controlada necesitó un promedio de 4,98 días para madurar, mientras que la fruta con SmartFresh<sup>SM</sup> requirió 5,44 días. A pesar de una variación considerable entre las muestras, algunas de las combinaciones cultivar/calibre presentaron patrones de maduración muy similares, ya sea en tratamiento con SmartFresh<sup>SM</sup> o almacenamiento en AC. En ambos cultivares, la fruta de mayor calibre con SmartFresh<sup>SM</sup> maduró más rápidamente que la fruta almacenada en AC, mientras que se verificó lo opuesto para los calibres más pequeños. Los resultados confirmaron nuestra recomendación previa en cuanto al contenido de

humedad del 70 % (30 % de masa seca) como punto límite para cosechar fruta, que se someta a tratamiento con SmartFresh<sup>SM</sup> para destinarse al sector del mercado “Maduro y listo para consumir” (en inglés, *Ripe and Ready*). Sin embargo, es posible enviar frutos de calibres de mayor tamaño (>300g) antes de alcanzar este punto, mientras que puede ser aconsejable no enviar los de calibres muy pequeños (<150g) a maduradores. Aunque la incidencia de la pulpa gris (decoloración difusa del mesocarpio) fue baja en el presente ensayo, por medio de un ensayo comercial realizado durante la estación anterior se demostró con claridad la capacidad de SmartFresh<sup>SM</sup> (1-metilciclopropeno) para reducir el desarrollo de la pulpa gris.

## **INTRODUCTION**

During 2000/1, the Agricultural Research Council's Institute for Tropical and Subtropical Crops launched a number of laboratory based trials with the ethylene inhibitor 1-methylcyclopropene (Lemmer & Kruger 2003)). This was followed by semi-commercial trials in static containers at Westfalia Estates during 2003 (Lemmer et al 2003). The first commercial 1-methylcyclopropene (1-MCP; Smartfresh<sup>SM</sup>) based exports took place during 2003. The primary aim of the application is to prevent soft landings and reduce physiological disorders such as greypulp (diffuse mesocarp discoloration) and pulp spot. Up to then, cold storage combined with controlled atmosphere (CA) was used for this purpose.

The first season's Smartfesh<sup>SM</sup> (SF) consignments all landed hard and were well received in Europe. However, certain English ripeners complained that the avocados may take too long to ripen and may ripen unevenly under certain conditions. A number of research trials were therefore initiated to address this aspect. The present paper deals with two of these. The first part concerns a laboratory trial that was conducted to determine an appropriate maturity threshold value for the 'Ripe and Ready' market sector while the second involves a commercial trial to confirm the latter recommendation under commercial conditions.

## **MATERIAL AND METHODS**

### **Laboratory based trials**

During 2004, SF was applied to 52 Hass samples from the Burgershall production region ranging in moisture content (MC) from 76% to 62% (24-38% dry mass content). The fruit were stored under export simulation conditions and the mean number of days to ripen (DTR) determined. This was compared with that of untreated fruit stored at regular atmosphere (RA).

## Commercial trial

During the second week of July 2006, SF was applied to 22 pallets of avocados destined for England. The container contained both Hass and Fuerte fruit originating from 5 farms managed by Westfalia Estates. The maturity levels of the samples varied between 71% and 62%. Upon arrival of the container in the United Kingdom during the first week of August, the content was split between Greencell (GC) in Spalding and Minor Weir & Willis (MWW) in Birmingham. Ripening was subsequently performed under commercial conditions. The evaluations entailed recording the DTR as well as scoring all physiological (lenticel damage, black cold injury, greypulp and bruising) and pathological (anthracnose, stem end rot and vascular browning) disorders.

## RESULTS AND DISCUSSION

The DTR results of the laboratory trial are visually displayed in Figure 1. At the beginning of the season, RA fruit with a MC between 76 and 75% required around 10 -11 DTR. At the same maturity level, SF fruit took approximately 12-13 DTR (2 days longer). The mean DTR of both RA and SF fruit steadily decreased as the season progressed. The DTR of both RA and SF stabilized at a MC of around 70%. Between 70% and 64%, the mean DTR of the RA fruit remained around 7 days. During the same period, the mean DTR of SF fruit was around 8 days.

The incidence of greypulp in Hass avocados from a selection of susceptible farms is shown in Figure 2. The disorder was found to become more prevalent as the season progressed but SF was found to reduce the incidence of greypulp in predisposed fruit.

The results of the commercial trial are summarized in Table 1. A total of 19 samples originating from 5 farms were ripened. In order to simplify the interpretation of the results, the same count originating from different farms are pooled in the table, resulting in ten sets of results. Four of these were of fruit ripened at GC (Fuerte count 14, Hass counts 14, 16 & 18) while the remaining six samples were ripened at MWW (Fuerte counts 12 & 16 and Hass counts 20, 22, 24 & 32).

The mean DTR of all CA fruit was 4.98 days whilst that of the SF fruit was 5.44 days. On average, the SF fruit thus took approximately half a day longer to ripen than the CA fruit, but the difference between the two means was not statistically different. The STD of the SF fruit was also half a day wider ( $\pm 1.8$  days) than that of the CA samples ( $\pm 1.26$ ), but the difference between the two mean STD's was not statistically significant either. Fruit count would seem to have influenced the differential ripening rate of the two treatments. In both Fuerte and Hass, the biggest SF fruit (Fuerte count 12 and Hass count 14) ripened slightly faster than their CA counterparts. In contrast, the smallest CA fruit (Hass 32) ripened faster

than the SF fruit. Considerable variation occurred with regard to the intermediate counts, but it would appear that, on average, the CA fruit ripened approximately half a day faster than the SF fruit. There did not seem to be a relationship between the mean DTR (both CA and SF) and the MC of the avocados at harvest. Neither would there seem to be a relationship between the STD of the MC on the one hand and the STD of the DTR on the other.

The ripening patterns of the ten samples are displayed in Figure 3 a – j. In count 12 Fuerte (Figure 3a), the SF peaked on day 5 when more than 50 % of the sample ripened. The CA peaked on day 4 when slightly less than 30 % of the sample ripened. All SF fruit were ripe by day 5 while around 10 % of CA fruit ripened on, respectively, days 6, 7 and 8. In the case of Fuerte count 14 (Figure 3b) the ripening patterns of the two samples were fairly similar in that around 45 % of both samples ripened on day 5. In the case of CA, more fruit ripened during the three days preceding the peak while the SF treatment had a longer tail. With count 16 Fuerte (Figure 3c) both treatments peaked on days 4 and 5. However, the CA fruit yielded between 40 and 45 % on these days while the SF fruit yielded between 25 and 30 % during this period and another 10 % during days 6 - 8. In the case of count 14 Hass (Figure 3d) the trend was opposite to the above. Both treatments again peaked on day 5 but the SF yielded between 35 – 40 % while the CA treatment yielded between 20 – 25 %. The CA fruit displayed a second smaller peak on days 9 and 10 during which approximately 10 % of the fruit ripened per day. The pattern was again reversed with Hass count 16 (Figure 3e). In this case, the ripening peak occurred on day 4 when nearly 40 % of CA fruit and close to 25 % of the SF fruit ripened. Although SF yielded more than CA on days 2 and 3, a second flatter peak occurred during days 9 to 12. In Hass count 18 (Figure 3f), both the CA and SF fruit peaked on day 5 when around 20 % of SF fruit and 25 % of CA fruit were ready to eat. Both treatments had a relatively long tail with the SF treatment having an additional smaller peak around day 10. Both CA and SF count 20 Hass avocados (Figure 4g) displayed a flat but distinct peak during days 3, 4, 5 and 6 when all fruit ripened (between 20 and 30 % per day) and no tail was present. With Hass count 22 (Figure 3h) a distinct peak occurred on day 5 when approximately 90 % of the CA fruit ripened. In the case of SF, between 20 and 30 % of the fruit ripened on day 3 and between 50 and 60 % on day 6. All count 24 CA Hass fruit (Figure 3i) ripened between days 3 and 5 while slightly less than 40 % of the SF fruit ripened on day 3 followed by slightly more than 20% on days 4 and 6. In Hass count 32 (Figure 3j) both treatments had two peaks. In the case of CA, the first peak of about 30 % occurred on day 3 followed by a second peak on day 5 when slightly more than 50% of the fruit ripened. In the case of SF, the first peak on day 3 was smaller when only 10 % of the fruit ripened with the bulk of fruit, around 80%, ripening during days 5 to 7.

The incidences of physiological and pathological disorders were low and no significant differences occurred between the two commercial treatments.

## **CONCLUSION**

The results indicate that, although considerable variation occurred between fruit, certain of the samples displayed very similar ripening patterns, whether treated with SF or stored under CA. In both cultivars, the largest SF fruit ripened faster than the CA fruit while the reverse was true for the smallest avocados. It would appear that the 70% moisture content cut off point for SF fruit destined for the 'Ripe and Ready' market sector is valid. Although the incidences of physiological and pathological disorders were low during the commercial trial, SF was found to significantly reduce the incidence of greypulp during the laboratory based trials.

## **ACKNOWLEDGEMENTS**

We sincerely thank Matthew Churchill of Greencell and Gerald Eva of Minor Weir and Willis, as well as their staff, for the efficient and professional manner in which the evaluations were conducted. We are also indebted to Ryan Larkan of Westfalia Marketing (UK) as well as to Jurg Bezuidenhout and the staff of Westfalia Packhouse for their kind cooperation. We would also like to thank Wehan Groenewald of Agrofresh, Rohm and Haas South Africa for financial support.

## REFERENCES

Lemmer, D. & Kruger, F.J. 2003. Laboratory based evaluation of 1-methyl cyclopropene (1-MCP) with five South African commercial export avocado cultivars. *Proceedings of the Fifth International World Avocado Congress*: 611-616.

Lemmer, D. Bezuidenhout, J., Sekhune. S., Ramokone, P, Letsoalo, L., Malumane, T.R, Chibi, P, Nxundu, Y. & Kruger, F.J. 2003. Semi-commercial evaluation of SmartFresh<sup>SM</sup> with South African export avocados in static containers at a packinghouse during 2002. *Proceedings of the Fifth International World Avocado Congress*: 617-622.

Table 1: Mean number of days required to ripen 3 samples of Fuerte and 7 samples of Hass avocados at two UK ripeners during August 2006.

Cultivar	Count	No of farms	N fruit CA/SF	Ripe-ner	Maturity (% moist)	Days to ripen	
						CA	SmartFresh
Fuerte	12	1	24/24	MWW	63.9 ± 1.6	5.0 ± 1.4	4.8 ± 1.6
Fuerte	14	2	56/56	GC	65.6 ± 2.6	4.2 ± 0.8	5.1 ± 1.3
Fuerte	16	1	47/45	MWW	63.3 ± 3.3	4.3 ± 0.5	5.0 ± 1.5
Hass	14	5	140/140	GC	NS	6.0 ± 2.5	5.2 ± 1.5
Hass	16	4	128/96	GC	67.7 ± 2.0	6.2 ± 2.0	7.0 ± 3.7
Hass	18	2	72/72	GC	67.3 ± 1.8	6.5 ± 2.1	7.5 ± 3.1
Hass	20	JP	34/30	MWW	71.5 ± 2.5	4.4 ± 1.1	4.4 ± 1.2
Hass	22	JP	35/30	MWW	70.1 ± 1.7	4.8 ± 0.5	5.1 ± 1.2
Hass	24	JP	47/43	MWW	68.3 ± 1.7	4.1 ± 0.8	4.6 ± 1.6
Hass	32	JP	70/75	MWW	67.5 ± 3.9	4.3 ± 0.9	5.7 ± 1.3
				Mean DTR		4.98 a	5.44 a
				Mean STD		1.26 a	1.80 a

CA : controlled atmosphere, SF: SmartFresh<sup>SM</sup>, MWW: Minor Weir and Willis, GC: Greencell, JP: jumble pack, DTR: days to ripen, STD: standard deviation




Figure 1: Mean number of days required to ripen Regular Atmosphere and SmartFresh<sup>SM</sup> Hass avocados at different maturity stages.




Figure 2: Incidence of geypulp in Regular Atmosphere and SmartFresh<sup>SM</sup> Hass avocados from specific farms after storage under export simulation conditions




Figure 3: Ripening patterns of 3 samples of Fuerte and 7 samples of Hass avocados at two UK ripeners during August 2006.